arXiv:1406.2662v1 [hep-th] 31 May 2014

On the Masses of the Universal hypermultiplets in heterotic M-theory

Nasr Ahmed

Astronomy Department, National Research Institute of Astronomy And Geophysics, Helwan, Cairo, Egypt.

1 abstract

The reduced 5D Heterotic M-theory has a deeply rich structure. For every Calabi-yau compactification, there exists a gravitational hypermultiplet $(g_{\mu\nu}, \psi_\mu, A_\mu)$ and a universal hypermultiplet. In this paper we derive the formulae for the masses of the scalar sector of the universal hypermultiplet $(V, \sigma, \zeta, \bar{\zeta})$ in the framework of 5D Heterotic M-theory.

2 Introduction

In the original formulation of M-theory [1, 2], all the standard model fields are trapped on two 9-branes located at the end points of an S^1/Z_2 orbifold. The 6 extra dimensions on the branes are compactified. A 5 dimensional reduction of the original HoravaWitten theory and the corresponding braneworld cosmology is given in [3, 4, 5].

In the 11-dimensional theory, the supergravity multiplet consists of the graviton field or the metric g, gravitino field ψ_I and a three index gauge field C_{IJK} with a field strength G_{IJKL}. The total bulk field content of this 5 dimensional theory is given by the gravity multiplet $(g_{\alpha\beta}, A_\alpha, \psi^i_\alpha)$ together with the universal hypermultiplet $(V, \sigma, \zeta, \bar{\zeta})$. A_α is a gauge field with field strength $\bar{F}_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$, ζ is a background complex field, V is the Calabi-Yau volume and ψ^i_α is the gravitino field. After the dualization, the three-form $C_{\alpha\beta\gamma}$ produces a scalar field σ. The 5 dimensional effective action can be written as [4]

$$S_5 = S_{bulk} + S_{bound} \tag{1}$$

where

$$\begin{aligned} S_{bulk} = \frac{-1}{2\kappa_5^2}\int_{\mathcal{M}_5}\sqrt{-g}\Big[\mathcal{R} + \frac{3}{2}\bar{F}_{\alpha\beta}\bar{F}^{\alpha\beta} \\ + \frac{1}{\sqrt{2}}\epsilon^{\alpha\beta\gamma\delta\epsilon}A_\alpha\bar{F}_{\beta\gamma}\bar{F}_{\delta\epsilon} + \frac{1}{2V^2}\partial_\alpha V\partial^\alpha V \\ + \frac{1}{2V^2}\left[(\partial_\alpha\sigma - i(\zeta\partial_\alpha\bar{\zeta} - \bar{\zeta}\partial_\alpha\zeta)\right. \\ \left. -2\alpha\epsilon(x^{11})A_\alpha)\right]^2 + \frac{2}{V}\partial_\alpha\zeta\partial^\alpha\bar{\zeta} + \frac{\alpha^2}{3V^2}\Big] \end{aligned} \tag{2}$$

and

$$S_{bound} = \frac{\sqrt{2}}{\kappa_5^2}\left[\int_{M_4^{(1)}}\sqrt{-g}V^{-1}\alpha - \int_{M_4^{(2)}}\sqrt{-g}V^{-1}\alpha\right] - \frac{1}{16\pi\alpha_{GUT}} \sum_{i=1}^{2}\int_{M_4^{(i)}}\sqrt{-g}\left(V\mathrm{tr}F_{\mu\nu}^{(i)}F^{(i)\mu\nu} - \sigma\mathrm{tr}F_{\mu\nu}^{(i)}\widetilde{F}^{(i)\mu\nu}\right). \quad (3)$$

where $\widetilde{F}^{(i)\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}^{(i)}$ and the expansion coeffecients α_i are

$$\alpha_i = \frac{\pi}{\sqrt{2}}\left(\frac{\kappa}{4\pi}\right)^{2/3}\frac{1}{v^{2/3}}\beta_i, \quad (4)$$
$$\beta_i = -\frac{1}{8\pi^2}\int_{C_i} tr(\mathcal{R}\wedge\mathcal{R}).$$

with the Calabi-Yau volume V defined as

$$V = \frac{1}{v}\int_X \sqrt{g^{(6)}} \quad (5)$$

where $g^{(6)}$ is the determinant of the Calabi-Yau metric.

3 The Non-linear Sigma model Lagrangian for the background V and ζ fields

In gaugino condensates the gaugino acquires non-zero vacuum expectation value which breaks the supersymmetry. The gaugino condensates lead to a background ζ field. So, now we have a background V field (represents the size of Calabi-Yau space), a background ζ field and a background metric. We take the following nonlinear sigma model lagrangian

$$L = \frac{-(\partial V)^2 - \left(\partial\sigma - i(\zeta\partial\bar{\zeta} - \bar{\zeta}\partial\zeta)\right)^2}{4V^2} - \frac{1}{V}(\partial\zeta)(\partial\bar{\zeta}) - U \quad (6)$$

Where $U = \frac{\alpha^2}{6V^2}$. We define the metric as

$$dh^2 = \frac{dV^2}{4V^2} + \frac{\left(d\sigma - i(\zeta d\bar{\zeta} - \bar{\zeta}d\zeta)\right)^2}{4V^2} + \frac{d\zeta d\bar{\zeta}}{V} \quad (7)$$

and introduce the one-forms u and v with their complex conjugate $\bar{u}$ and $\bar{v}$

$$w^a = (u, \bar{u}, v, \bar{v}) \quad (8)$$

Where

$$u = \frac{d\zeta}{\sqrt{V}} \quad (9)$$

And

$$v = \frac{1}{2V}\left(dV + id\sigma + \zeta d\bar{\zeta} - \bar{\zeta}d\zeta\right) \quad (10)$$

This leads to

$$du = -\frac{1}{2}(v+\bar{v})\wedge u \quad (11)$$

$$dv = -\bar{v}\wedge v - \bar{u}\wedge u \quad (12)$$

and the connection two-forms

$$w^v{}_v = \bar{v} - v \quad (13)$$
$$w^u{}_u = \frac{1}{2}(\bar{v} - v) \quad (14)$$
$$w^u{}_v = -u \quad (15)$$

Cartan's structure equantions are

$$T^a = d\theta^a + w_b^a\wedge\theta^b \quad (16)$$
$$\Omega_b^a = dw_b^a + w_c^a\wedge w_b^c \quad (17)$$

For the torsion and curvature 2-form respectively.
The curvature 2-form gives (when expressed locally)

$$\Omega^a_b = \frac{1}{2} R^a_{bcd} \theta^c \wedge \theta^d \tag{18}$$

Where the components R^a_{bcd} are in orthonormal basis. The affine connection form satisfies

$$w_{ab} + w_{ba} = dg_{ab} \quad , \quad w_{ab} = g_{ac} w^c_b \tag{19}$$

Since for the orthonormal metric $g_{ab} = \eta_{ab}$ is constant, we have For the curvature 2-form we have

$$\Omega_{ab} = g_{ac} \Omega^c_b = -\Omega_{ba} \tag{20}$$

For a free torsion space, Cartan's first structure equation is

$$d\theta^a = -w^a_b \wedge \theta^b \tag{21}$$

So, the key formulas we are going to use to derive the connection coefficients and the corresponding curvature tensor are

$$\begin{aligned} w_{ab} &= -w_{ba} && (22)\\ d\theta^a &= -w^a_b \wedge \theta^b && (23)\\ \frac{1}{2} R^a_{bcd} \theta^c \wedge \theta^d &= dw^a_b + w^a_c \wedge w^c_b && (24) \end{aligned}$$

For the Ricci tensor we find

$$\begin{aligned} R^u{}_u &= \frac{1}{2} R^u{}_{u\alpha\beta} w^\alpha \wedge w^\beta \\ &= (\bar{v} \wedge v) + (\bar{u} \wedge u) \end{aligned} \tag{25}$$

And

$$R^v{}_v = (\bar{u} \wedge u) - (\bar{v} \wedge v) \tag{26}$$

The components of the curvature tensor are ($g_{u\bar{u}} = \frac{1}{2}$):

$$\begin{gathered} R_{u\bar{u}u\bar{u}} = 1, \quad R_{u\bar{u}v\bar{v}} = \frac{1}{2}, \quad R_{v\bar{v}v\bar{v}} = 1, \\ R_{u\bar{v}v\bar{u}} = \frac{1}{2}, \quad R_{\bar{u}uu\bar{u}} = -1, \quad R_{u\bar{u}\bar{u}u} = -1, \\ R_{\bar{u}uv\bar{v}} = -\frac{1}{2}, \; R_{u\bar{u}\bar{v}v} = -\frac{1}{2}, \quad R_{\bar{v}uv\bar{u}} = -\frac{1}{2}, \\ R_{\bar{v}vv\bar{v}} = -1, \quad R_{u\bar{v}\bar{u}v} = -\frac{1}{2}, \quad R_{v\bar{u}u\bar{v}} = \frac{1}{2}, \\ R_{\bar{u}u\bar{u}u} = 1, \quad R_{\bar{v}v\bar{v}v} = 1, \quad R_{\bar{u}u\bar{v}v} = \frac{1}{2}, \\ R_{\bar{v}vu\bar{u}} = -\frac{1}{2}, \quad R_{v\bar{v}\bar{u}u} = -\frac{1}{2}, \quad R_{\bar{v}v\bar{u}u} = \frac{1}{2}, \\ R_{\bar{v}u\bar{u}v} = \frac{1}{2}, \quad R_{\bar{u}vu\bar{v}} = \frac{1}{2}, \quad R_{v\bar{u}\bar{v}u} = -\frac{1}{2}, \\ R_{\bar{u}v\bar{v}u} = \frac{1}{2}, \quad R_{v\bar{v}u\bar{u}} = \frac{1}{2} R_{v\bar{v}\bar{v}v} = -1. \end{gathered}$$

The grad of the potential U is

$$\nabla U = -\frac{\alpha^2}{3V^2}(v + \bar{v}) \tag{27}$$

$$\begin{aligned} \nabla^2 U = \frac{2\alpha^2}{3V^2}(v + \bar{v}) \otimes (v + \bar{v}) - \frac{\alpha^2}{3V^2} \\ (v - \bar{v}) \otimes (v - \bar{v}) + \frac{2\alpha^2}{3V^2} u \otimes \bar{u} \end{aligned} \tag{28}$$

Now we would like to express the the lagrangian (7) in terms of the one-forms u and v. We make use of the general form of the nonlinear sigma model lagrangian with a background field ζ

$$\begin{aligned} L &= -\frac{1}{2} g_{ij} (D_\mu \zeta^i)(D^\mu \zeta^j) \\ &\quad + \frac{1}{2} (\partial_\mu \phi^i)(\partial^\mu \phi^j) R_{ikjl} \zeta^k \zeta^l - \frac{1}{2} U_{;ij} \zeta^i \zeta^j. \end{aligned} \tag{29}$$

Where R_{ijkl} is the curvature of g_{ij}. After some manipulations we get the lagrangian in the form

$$L = L_1 + L_2 \tag{30}$$

Where

$$L_1 = -(D_\alpha \zeta^u)(D^\alpha \zeta^{\bar{u}}) - (D_\alpha \zeta^v)(D^\alpha \zeta^{\bar{v}}) \quad (31)$$

And

$$\begin{aligned} L_2 = & \frac{\alpha^2}{2} V^{-2} \left[(\zeta^v - \zeta^{\bar{v}}) + C(\zeta^u - \zeta^{\bar{u}}) \right]^2 \quad (32) \\ & - \frac{\alpha^2}{2} V^{-2} (\zeta^u - C\zeta^v)(\zeta^{\bar{u}} - C\zeta^{\bar{v}}) \\ & - \frac{\alpha^2}{2} V^{-2} (\zeta^v + \zeta^{\bar{v}})^2 - \frac{\alpha^2}{3} V^{-2} \zeta^u \zeta^{\bar{u}} \end{aligned}$$

The first part L_1 is diagonalized and we need to diagonalize the second part L_2 to get the eigen Values. To do that we make the following change of variables

$$\zeta^v = \frac{1}{\sqrt{2}}(X + iY) \quad , \quad \zeta^u = \frac{1}{\sqrt{2}}(Z + iW) \quad (33)$$

That means we have 4 fields X,Y,Z, and W. In terms of the new fields, the lagrangian L_2 becomes

$$\begin{aligned} L_2 &= -\alpha^2 V^{-2} (Y + CW)^2 \quad (34) \\ &- \frac{\alpha^2}{4} V^{-2} \left((Z - CX)^2 + (W - CY)^2 \right) \\ &- \frac{\alpha^2}{3} V^{-2} \left(2X^2 + Y^2 + \frac{1}{2} Z^2 + \frac{1}{2} W^2 \right). \end{aligned}$$

Which could be written in a matrix form as

$$L_2 = -\frac{\alpha^2}{V^2} \begin{pmatrix} \frac{2}{3} + \frac{C^2}{4} & 0 & -\frac{C}{2} & 0 \\ 0 & \frac{4}{3} + \frac{C^2}{4} & 0 & 2C \\ -\frac{C}{2} & 0 & \frac{5}{12} & 0 \\ 0 & -\frac{C}{2} & 0 & \frac{5}{12} + C^2 \end{pmatrix}$$

After diagonalization , The eigen values that represents the masses of the scalar sector of the universal hypermultiplet $(V, \sigma, \zeta, \bar{\zeta})$ are

$$\frac{\alpha^2}{24V^2} \left(-21 - 15C^2 + \sqrt{121 - 774C^2 + 81C^4} \right), \quad (35)$$

$$\frac{\alpha^2}{24V^2} \left(-21 - 15C^2 + \sqrt{121 - 774C^2 + 81C^4} \right), \quad (36)$$

$$\frac{\alpha^2}{24V^2} \left(-13 - 3C^2 + 3\sqrt{1 + 18C^2 + C^4} \right), \quad (37)$$

$$\frac{\alpha^2}{24V^2} \left(-13 - 3C^2 - 3\sqrt{1 + 18C^2 + C^4} \right). \quad (38)$$

4 Conclusion

Making use of the non-linear sigma model Lagrangian of the background fields, We derived the formulae for the masses of the scalar sector of the universal hypermultiplet $(V, \sigma, \zeta, \bar{\zeta})$ in the reduced 5D Heterotic M-theory.

5 Acknwoledgment

I am so grateful to Prof. Ian Moss from Newcastle university for the very useful discussions during this work.

References

[1] Horava, P., and Witten, E. Heterotic and Type I string dynamics from eleven dimensions. Nucl. Phys. B 460 (1996) 506524 [hep-th/9510209].

[2] Horava, P., and Witten, E. Eleven-Dimensional Supergravity on a Manifold with Boundary. Nucl. Phys. B475 (1996) 94-114 [hep-th/9603142].

[3] Lukas, A., Ovrut, B.A., Stelle, K.S., and Waldram, D. The Universe as a domain wall. Phys. Rev. D, 59 (1999) 08600119 [hep-th/9803235].

[4] Lukas, A., Ovrut, B.A., and Waldram, D. Cosmological solutions of HoravaWitten theory. Phys. Rev. D, 60 (1999) 086001111 [hep-th/9806022].

[5] Lukas, A., Ovrut, B.A., and Waldram, D. Boundary inflation. Phys. Rev. D, 61 (2000) 023506 118.

[6] Z. Lalak and S. Thomas. Gaugino Condensation, Moduli Potential and Supersymmetry Breaking in $M-$theory Models. Nucl. Phys. B515 (1998) 55-72 [hep-th/9707223].